

Rational Calogero models based on rank-2 root systems: supertraces on the superalgebras of observables

S.E.Konstein ^{*†}

*I.E.Tamm Department of Theoretical Physics,
P. N. Lebedev Physical Institute,
117924, Leninsky Prospect 53, Moscow, Russia.*

Abstract

It is shown that the superalgebra $H_{W(I_2(n))}$ of observables of the rational Calogero model based on the root system $I_2(n)$ possesses $[(n+1)/2]$ supertraces. Model with three-particle interaction based on the root system G_2 belongs to this class of models and its superalgebra of observables has 3 independent supertraces.

^{*}E-mail: konstein@td.lpi.ac.ru

[†]This work is supported by the Russian Fund for Basic Research, Grants 96-02-17314 and 96-15-96463.

1 Introduction

In this work we continue to compute the number of supertraces on the superalgebras of observables underlying the rational Calogero models [1] based on the root systems [2]. The results for the root systems of A_{N-1} , B_N (C_N) and D_N types are listed in [3], [4]. Here we consider the root systems of $I_2(n)$ type. As $W(A_2) = W(I_2(3))$, $W(B_2) = W(I_2(4))$, $W(D_2) = W(I_2(2))$ and $W(G_2) = W(I_2(6))$ the case under consideration covers all rank-2 systems. It is shown that the number of supertraces on the superalgebra of observables of Calogero model based on this root system is $[(n+1)/2]$. In particular it gives that in the case of the root system G_2 there are 3 supertraces.

The definition and some properties of the superalgebra of observables are discussed in the next section. In the section 3 the condition sufficient for existence of the supertraces is formulated and some consequences from the existence of the supertraces are discussed. The superalgebra $H_{W(I_2(n))}$ of observables of the rational Calogero model based on the root system $I_2(n)$ is described in the section 4 and the number of supertraces on this superalgebra is computed in the last section.

2 The superalgebra of observables

The superalgebra of observables of the rational Calogero model based on the root system is defined in the following way.

Let us define the reflections $R_{\vec{v}} \forall \vec{v} \in \mathbb{R}_N$, $\vec{v} \neq 0$, as follows

$$R_{\vec{v}}(\vec{x}) = \vec{x} - 2 \frac{(\vec{x}, \vec{v})}{(\vec{v}, \vec{v})} \vec{v} \quad \forall \vec{x} \in \mathbb{R}_N. \quad (1)$$

Here the notation $(\cdot, \cdot)$ used for the inner product in $\mathbb{R}_N$. We will use also the coordinates of vectors, $v_i \stackrel{def}{=} (\vec{v}, \vec{e}_i)$, where vectors $\vec{e}_i$ constitute the orthonormal basis in $\mathbb{R}_N$: $(\vec{e}_i, \vec{e}_j) = \delta_{ij}$. The reflections (1) satisfy the properties

$$R_{\vec{v}}(\vec{v}) = -\vec{v}, \quad R_{\vec{v}}^2 = 1, \quad (R_{\vec{v}}(\vec{x}), \vec{u}) = (\vec{x}, R_{\vec{v}}(\vec{u})), \quad \forall \vec{v}, \vec{x}, \vec{u} \in \mathbb{R}_N. \quad (2)$$

The finite set of vectors $\mathbf{R} \subset \mathbb{R}_N$ is the *root system* if $\mathbf{R}$ is $R_{\vec{v}}$ -invariant $\forall \vec{v} \in \mathbf{R}$ and the group $W(\mathbf{R})$ generated by all reflections $R_{\vec{v}}$ with $\vec{v} \in \mathbf{R}$ (Coxeter group) is finite. The brief description of the classification of the root systems one can find for instance in [2].

The group $W(\mathbf{R})$ acts also on some space $\mathbf{C}$ of functions on $\mathbb{R}_N$. Let us assume for definiteness that $\mathbf{C}$ is constituted by all infinitely smooth functions of polynomial growth *i.e.* that $\forall f \in \mathbf{C} \exists n(f) \in \mathbb{Z}_+$ such that $\lim_{\vec{x} \rightarrow \infty} f(\vec{x})/|\vec{x}|^{n(f)} = 0$. Every function $f \in \mathbf{C}$ can be also considered as element of $End \mathbf{C}$ *i.e.* as operator on $\mathbf{C}$ acting by multiplication: $g \rightarrow fg \forall g \in \mathbf{C}$. The action of $R_{\vec{v}} \in W(\mathbf{R})$ on such functions has the following form

$$R_{\vec{v}}(f(\vec{x})) = f(R_{\vec{v}}(\vec{x})) \text{ when } f(\vec{x}) \in \mathbf{C}, \quad (3)$$

$$R_{\vec{v}}f(\vec{x}) = f(R_{\vec{v}}(\vec{x}))R_{\vec{v}} \text{ when } f(\vec{x}) \in End \mathbf{C}. \quad (4)$$

Dunkl differential-difference operators is defined as [5]

$$D_i = \frac{\partial}{\partial x_i} + \sum_{\vec{v} \in \mathbf{R}} \nu_{\vec{v}} \frac{v_i}{(\vec{x}, \vec{v})} (1 - R_{\vec{v}}), \quad (5)$$

where coupling constants $\nu_{\vec{v}}$ are such that

$$\nu_{\vec{v}} = \nu_{R_{\vec{u}}(\vec{v})} \quad \forall \vec{u}, \vec{v} \in \mathbf{R}. \quad (6)$$

These operators are well defined on $\mathbf{C}$ and commute with each other, $[D_i, D_j] = 0$.

Due to this property the deformed creation and annihilation operators [6, 7]

$$a_i^\alpha = \frac{1}{\sqrt{2}}(x_i + (-1)^\alpha D_i), \quad \alpha = 0, 1, \quad (7)$$

transform under action of reflections $R_{\vec{v}} \quad \forall \vec{v} \in \mathbf{R}$ as $\mathbb{R}_N$ vectors

$$R_{\vec{v}} a_i^\alpha = \sum_{j=1}^N \left(\delta_{ij} - 2 \frac{v_i v_j}{(\vec{v}, \vec{v})} \right) a_j^\alpha R_{\vec{v}} \quad (8)$$

and satisfy the following commutation relations

$$[a_i^\alpha, a_j^\beta] = \varepsilon^{\alpha\beta} \left(\delta_{ij} + 2 \sum_{\vec{v} \in \mathbf{R}} \nu_{\vec{v}} \frac{v_i v_j}{(\vec{v}, \vec{v})} R_{\vec{v}} \right) \quad (9)$$

where $\varepsilon^{\alpha\beta}$ is the antisymmetric tensor, $\varepsilon^{01} = 1$.

The operators a_i^α ($\alpha = 0, 1, i = 1, \dots, N$) together with the elements of the group $W(\mathbf{R})$ generate the associative algebra with elements polynomial on a_i^α . We denote this algebra as $H_{W(\mathbf{R})}(\nu)$ and call it *the algebra of observables of Calogero model based on the root system $\mathbf{R}$* . Here the notation ν stands for a complete set of $\nu_{\vec{v}}$ with $\vec{v} \in \mathbf{R}$.

The commutation relations (9) and (8) allows one to define the parity π :

$$\pi(a_i^\alpha) = 1 \quad \forall \alpha, i, \quad \pi(g) = 0 \quad \forall g \in W(\mathbf{R}) \quad (10)$$

and consider $H_{W(\mathbf{R})}(\nu)$ as a superalgebra.

Obviously the superalgebra $H_{W(\mathbf{R})}(\nu)$ contains as a subalgebra the group algebra $\mathbb{C}[W(\mathbf{R})]$ of Coxeter group $W(\mathbf{R})$.

An important property of superalgebra $H_{W(\mathbf{R})}(\nu)$ is that it has $\mathfrak{sl}_2$ algebra of inner differentiations with the generators

$$T^{\alpha\beta} = \frac{1}{2} \sum_{i=1}^N \{a_i^\alpha, a_i^\beta\} \quad (11)$$

which commute with $\mathbb{C}[W(\mathbf{R})]$, $[T^{\alpha\beta}, R_{\vec{v}}] = 0$, and act on a_i^α as on $\mathfrak{sl}_2$ -vectors:

$$[T^{\alpha\beta}, a_i^\gamma] = \varepsilon^{\alpha\gamma} a_i^\beta + \varepsilon^{\beta\gamma} a_i^\alpha. \quad (12)$$

The restriction of operator $\hat{H}_{Cal}^{\mathbf{R}}(\nu) \stackrel{def}{=} T^{01}$ on the subspace $\mathbf{C}_{W(\mathbf{R})} \subset \mathbf{C}$ of $W(\mathbf{R})$ -invariant functions is the second-order differential operator which is well-known Hamiltonian of the rational Calogero model [1] based on the root system $\mathbf{R}$ [2]. One of the relations (11) namely $[\hat{H}_{Cal}^{\mathbf{R}}(\nu), a_i^\alpha] = -(-1)^\alpha a_i^\alpha$ allows one to find the wavefunctions of the equation $\hat{H}_{Cal}^{\mathbf{R}}(\nu)\psi = \epsilon\psi$ via usual Fock procedure with the vacuum $|0\rangle$ such that $a_i^0|0\rangle=0 \quad \forall i$ [7]. After $W(\mathbf{R})$ -symmetrization these wavefunctions become the wavefunctions of Calogero Hamiltonian $H_{Cal}^{\mathbf{R}}(\nu)|_{\mathbf{C}_{W(\mathbf{R})}}$.

3 Supertraces on $H_{W(\mathbf{R})}(\nu)$

Definition. The supertrace on the superalgebra $\mathcal{A}$ is a linear complex-valued function $str(\cdot)$ such that $\forall f, g \in \mathcal{A}$ with definite parity $\pi(f)$ and $\pi(g)$

$$str(fg) = (-1)^{\pi(f)\pi(g)} str(gf). \quad (13)$$

Every supertrace $str(\cdot)$ on $\mathcal{A}$ generates the invariant bilinear form on $\mathcal{A}$

$$B_{str}(f, g) = str(f \cdot g). \quad (14)$$

It is obvious that if such a bilinear form is degenerated then the null-vectors of this form constitute both-sided ideal $\mathcal{I} \subset \mathcal{A}$.

It was shown that the ideals of this sort are present in the superalgebras $H_{W(A_1)}(\nu)$ (corresponding to the two-particle Calogero model) at $\nu = k + \frac{1}{2}$ [8] and in the superalgebras $H_{W(A_2)}(\nu)$ (corresponding to three-particle Calogero model) at $\nu = k + \frac{1}{2}$ and $\nu = k \pm \frac{1}{3}$ [9] with every integer k and that for all the other values of ν all supertraces on these superalgebras generate the nondegenerated bilinear forms (14).

The spectrum of N -particle rational Calogero Hamiltonian (case $\mathbf{R} = A_{N-1}$) coincides with the spectrum of system of N noninteracting oscillators if the latter is shifted on the constant $c(\nu) = \frac{1}{2}N(N-1)\nu$. It allows one to construct the similarity transformation between operators $\hat{H}_{Cal}^{A_{N-1}}(\nu)|_{\mathbb{C}[W(\mathbf{R})]} - c(\nu)$ with different ν [10]. Nevertheless the previous consideration shows that the corresponding algebras $H_{W(A_{N-1})}(\nu)$ can be nonisomorphic at different values of ν .

It is easy to describe all supertraces on $\mathbb{C}[W(\mathbf{R})]$. Every supertrace on $\mathbb{C}[W(\mathbf{R})]$ is completely defined by its values on $W(\mathbf{R}) \subset \mathbb{C}[W(\mathbf{R})]$ and the function str is a central function on $W(\mathbf{R})$ i.e. the function on the conjugacy classes.

To formulate the theorem establishing the connection between the supertraces on $H_{W(A_N)}(\nu)$ and the supertraces on $\mathbb{C}[W(\mathbf{R})]$ let us introduce the grading on the vector space of $\mathbb{C}[W(\mathbf{R})]$. The grading E of elements $g \in W(\mathbf{R}) \subset \mathbb{C}[W(\mathbf{R})]$ is defined as follows. Let $\mathcal{H}_N^\alpha$ be the linear space with basis elements $a_1^\alpha, a_2^\alpha, \dots, a_N^\alpha$. Consider the subspaces $\mathcal{E}^\alpha(g) \subset \mathcal{H}_N^\alpha$ as

$$\mathcal{E}^\alpha(g) = \{h \in \mathcal{H}_N^\alpha : gh = -hg\}, \quad (15)$$

and put

$$E(g) = \dim \mathcal{E}^\alpha(g). \quad (16)$$

To avoid misunderstanding it should be noticed that $\mathbb{C}[W(\mathbf{R})]$ is not in general a graded algebra.

The following theorem was proved in [3]¹:

¹this theorem was proved for the case $\mathbf{R} = A_N$ only but the proof does not depend on the particular properties of the symmetric group $S_N = W(A_{N-1})$.

Theorem 1. Let $\mathcal{P}(g)$ be the projector $\mathbb{C}[W(\mathbf{R})] \rightarrow \mathbb{C}[W(\mathbf{R})]$ defined as $\mathcal{P}(\sum_i \alpha_i g_i) = \sum_{i: g_i \neq 1} \alpha_i g_i$ ($g_i \in W(\mathbf{R})$, $\alpha_i \in \mathbb{C}$). Let the grading E defined in (16) and the subspaces $\mathcal{E}^\alpha(g)$ defined in (15) satisfy the equations

$$E(\mathcal{P}([h_0, h_1])g) = E(g) - 1 \quad \forall g \in \mathbb{C}[W(\mathbf{R})], \quad \forall h_\alpha \in \mathcal{E}^\alpha(g). \quad (17)$$

Then every supertrace on the algebra $\mathbb{C}[W(\mathbf{R})]$ satisfying the equations

$$\text{str}([h_0, h_1]g) = 0 \quad \forall g \in W(\mathbf{R}) \text{ with } E(g) \neq 0 \text{ and } \forall h_\alpha \in \mathcal{E}^\alpha(g), \quad (18)$$

can be extended to the supertrace on $H_{W(\mathbf{R})}(\nu)$ in a unique way.

4 Superalgebras $H_{W(I_2(n))}$

It is convenient to use $\mathbb{C}$ instead of $\mathbb{R}_2$ to describe $H_{W(I_2(n))}$. The root system $I_2(n)$ contains $2n$ vectors $v_k = \exp(\pi i k/n)$, $k = 0, 1, \dots, 2n-1$. The corresponding Coxeter group $W(I_2(n))$ has $2n$ elements, n reflections R_k acting on z , $z^* \in \mathbb{C}$ as follows

$$\begin{aligned} R_k z &= -z^* v_k^2 R_k, \\ R_k z^* &= -z v_k^{*2} R_k, \end{aligned} \quad k \in \mathbb{Z}_n \quad (19)$$

and n elements of the form $S_k = R_k R_0$. S_0 is the unity in $W(I_2(n))$. These elements satisfy the following relations

$$R_k R_l = S_{k-l}, \quad S_k S_l = S_{k+l}, \quad R_k S_l = R_{k-l}, \quad S_k R_l = R_{k+l}. \quad (20)$$

Obviously the reflections R_{2k} lie in one conjugacy class and R_{2k+1} in another if n is even. If n is odd then all reflections R_k lie in one conjugacy class.

It is convenient to consider the following basis in $\mathbb{C}[W(I_2(n))]$

$$L_p = \sum_{k=0}^{n-1} \lambda^{kp} R_k, \quad Q_p = \sum_{k=0}^{n-1} \lambda^{-kp} S_k, \quad (21)$$

$$\text{where } \lambda = \exp(2\pi i/n), \quad p \in \mathbb{Z}_n. \quad (22)$$

Differential-difference operators have the following form

$$\begin{aligned} D_z &= 2 \frac{\partial}{\partial z^*} + \nu_0 \sum_{k=0}^{n-1} \frac{v_{2k}}{z v_{2k}^* + z^* v_{2k}} (1 - R_{2k}) + \nu_1 \sum_{k=0}^{n-1} \frac{v_{2k+1}}{z v_{2k+1}^* + z^* v_{2k+1}} (1 - R_{2k+1}), \\ D_{z^*} &= 2 \frac{\partial}{\partial z} + \nu_0 \sum_{k=0}^{n-1} \frac{v_{2k}^*}{z v_{2k}^* + z^* v_{2k}} (1 - R_{2k}) + \nu_1 \sum_{k=0}^{n-1} \frac{v_{2k+1}^*}{z v_{2k+1}^* + z^* v_{2k+1}} (1 - R_{2k+1}). \end{aligned} \quad (23)$$

This form unifies both cases of even and odd n . If n is odd then D_z and D_{z^*} depends on the only coupling constant $\nu_0 + \nu_1$.

The basis in the space $\mathcal{H}_2^\alpha$ is

$$\begin{aligned} a^\alpha &\stackrel{\text{def}}{=} \frac{1}{2} (z + (-1)^\alpha D_z) \\ b^\alpha &\stackrel{\text{def}}{=} \frac{1}{2} (z^* + (-1)^\alpha D_{z^*}). \end{aligned} \quad (24)$$

Now we can write down the relation between elements a^α , b^α , R_k and S_k generating the associative superalgebra $H_{W(I_2(n))}$ of polynomials of a_i^α , b_i^α with coefficients in $\mathbb{C}[W(I_2(n))]$:

$$R_k a^\alpha = -v_k^2 b^\alpha R_k, \quad R_k b^\alpha = -v_k^{*2} a^\alpha R_k, \quad (25)$$

$$S_k a^\alpha = v_k^{*2} a^\alpha S_k, \quad S_k b^\alpha = v_k^2 b^\alpha S_k, \quad (26)$$

$$L_p a^\alpha = -b^\alpha L_{p+1}, \quad L_p b^\alpha = -a^\alpha L_{p-1}, \quad (27)$$

$$Q_p a^\alpha = a^\alpha Q_{p+1}, \quad Q_p b^\alpha = b^\alpha Q_{p-1}, \quad (28)$$

$$L_k L_l = n \delta_{k+l} Q_l, \quad L_k Q_l = n \delta_{k-l} L_l, \quad (29)$$

$$Q_k L_l = n \delta_{k+l} L_l, \quad Q_k Q_l = n \delta_{k-l} Q_l, \quad \text{where } \delta_k \stackrel{\text{def}}{=} \delta_{k0}, \quad (29)$$

$$\begin{aligned} [a^\alpha, b^\beta] &= \varepsilon^{\alpha\beta} \left(1 + \frac{\nu_0 + \nu_1}{2} L_0 - \frac{\nu_0 - \nu_1}{2} L_{n/2} \right), \\ [a^\alpha, a^\beta] &= \varepsilon^{\alpha\beta} \left(\frac{\nu_0 + \nu_1}{2} L_1 - \frac{\nu_0 - \nu_1}{2} L_{n/2+1} \right), \\ [b^\alpha, b^\beta] &= \varepsilon^{\alpha\beta} \left(\frac{\nu_0 + \nu_1}{2} L_{-1} - \frac{\nu_0 - \nu_1}{2} L_{n/2-1} \right). \end{aligned} \quad (30)$$

The terms containing $\nu_0 - \nu_1$ in (30) are absent when n is odd and halfinteger indices are senseless, so let us assume that $\nu_0 = \nu_1$ when n is odd.

5 The number of supertraces on $H_{W(I_2(n))}$

In this section the following theorem is proved:

Theorem 2. *The superalgebra $H_{W(I_2(n))}$ has $\left\lfloor \frac{n+1}{2} \right\rfloor$ supertraces.*

It is easy to find the grading E :

$$\begin{aligned} E(R_k) &= 1, & R_k(v_k^* a^\alpha + v_k b^\alpha) &= -(v_k^* a^\alpha + v_k b^\alpha) R_k, \\ E(S_k) &= 0, & \forall k &\neq \frac{n}{2}, \\ E(S_{n/2}) &= 2, & S_{n/2} a^\alpha &= -a^\alpha S_{n/2}, \quad S_{n/2} b^\alpha = -b^\alpha S_{n/2}, \text{ if } n \text{ is even and } S_{n/2} \text{ exists,} \end{aligned} \quad (31)$$

and check that $H_{W(I_2(n))}$ satisfies the conditions of Theorem 1, i.e. that

$$E\left(\mathcal{P}\left([v_k^* a^0 + v_k b^0], [v_k^* a^1 + v_k b^1]\right) R_k\right) = 0 \quad (32)$$

and that for even n

$$\begin{aligned} E\left(\mathcal{P}\left([a^0, a^1]\right) S_{n/2}\right) &= E\left(\mathcal{P}\left([a^0, b^1]\right) S_{n/2}\right) = \\ E\left(\mathcal{P}\left([b^0, a^1]\right) S_{n/2}\right) &= E\left(\mathcal{P}\left([b^0, b^1]\right) S_{n/2}\right) = 1. \end{aligned} \quad (33)$$

To compute the number of supertraces on the superalgebra we have to find the number of the solutions of the equations (18) which have the following form for the algebra under consideration

$$\text{str}\left([v_k^* a^0 + v_k b^0], [v_k^* a^1 + v_k b^1]\right) R_k = 0 \quad (34)$$

and

$$\begin{aligned} \text{str} \left([a^0, a^1] S_{n/2} \right) &= \text{str} \left([a^0, b^1] S_{n/2} \right) = \\ \text{str} \left([b^0, a^1] S_{n/2} \right) &= \text{str} \left([b^0, b^1] S_{n/2} \right) = 0 \quad \text{for even } n. \end{aligned} \quad (35)$$

The equations (34) lead to

$$\begin{aligned} \text{str}(R_k) = & - \frac{\nu_0 + \nu_1}{2} \text{str} \left(Q_0 + \frac{1}{2}(Q_1 + Q_{-1}) \right) \\ & + (-1)^k \frac{\nu_0 - \nu_1}{2} \text{str} \left(Q_{n/2} + \frac{1}{2}(Q_{n/2+1} + Q_{n/2-1}) \right), \end{aligned} \quad (36)$$

and when n is even the relations (35) take place and lead to

$$\text{str} \left((\nu_0 + \nu_1) L_{\pm 1} - (-1)^{n/2} (\nu_0 - \nu_1) L_{n/2 \pm 1} \right) = 0 \quad (37)$$

$$\text{str}(S_{n/2}) = -\frac{1}{2} \text{str} \left((\nu_0 + \nu_1) L_0 - (-1)^{n/2} (\nu_0 - \nu_1) L_{n/2} \right). \quad (38)$$

It is easy to see that equations (37) are consequences of (36). The equations (36) express $\text{str}(R_k) \forall k$ via $\text{str}(S_j)$ and (38) expresses $\text{str}(S_{n/2})$ via $\text{str}(S_j)$ with $j \neq n/2$ when n is even. Hence due to theorem 1 every supertrace on $H_{W(I_1(n))}(\nu)$ is determined completely by its values on S_j with $j \neq n/2$. Since S_j and S_k belong to one conjugacy class if and only if $j = -k$ the number of independent supertrace is equal to $(n+1)/2$ when n is odd and to $n/2$ if n is even. It finishes the proof of Theorem 2.

For $n = 2, 3, 4$ the obtained result are in agreement with the results obtained in [3] and [4]. The case $n = 6$ gives that the superalgebra of observables of the rational Calogero model based on the root system of G_2 type has 3 independent supertraces.

References

- [1] F. Calogero, J. Math. Phys., **10** (1969) 2191, 2197; *ibid* **12** (1971) 419.
- [2] M. A. Olshanetsky and A. M. Perelomov, Phys. Rep., **94** (1983) 313.
- [3] S.E. Konstein and M.A. Vasiliev, J. Math. Phys. **37** (1996) 2872.
- [4] S.E. Konstein, Teor. Mat. Fiz., **111** (1997) 592.
- [5] C.F. Dunkl, Trans. Am. Math. Soc. **311** (1989) 167.
- [6] A. Polychronakos, Phys. Rev. Lett. **69** (1992) 703.
- [7] L. Brink, H. Hansson and M.A. Vasiliev, Phys. Lett. **B286** (1992) 109.
- [8] M.A. Vasiliev, JETP Letters, **50** (1989) 344-347; Int. J. Mod. Phys. **A6** (1991) 1115.
- [9] S.E. Konstein, preprint FIAN/TD/97-17.
- [10] N. Gurappa and P.K. Panigrahi, cond-mat/9710035.